

Mathematics Questions by Topic

Motion and Force

Answer 48 Source: K7SM2S21

Question 48

A light inextensible string passes over a smooth pulley. Particles of masses m_1 and m_2 are attached to each end of the string as shown in the diagram. If the mass m_2 accelerates

downwards at $\frac{g}{5} \text{ m/s}^2$, then the ratio $\frac{m_2}{m_1}$ is equal to

- A. 1
- B. $\frac{3}{2}$
- C. $\frac{2}{3}$
- D. 5
- E. $\frac{5}{4}$

ANSWER B

Consider the mass m_2 moving downwards, resolving downwards,

$$(1) \quad m_2 g - T = m_2 a$$

Consider the mass m_1 moving upwards, resolving upwards,

$$(2) \quad T - m_1 g = m_1 a$$

to solve for a add the two equations to eliminate T

$$m_2 g - m_1 g = m_2 a + m_1 a$$

$$\text{so that } (m_2 - m_1)g = (m_1 + m_2)a$$

$$a = \frac{(m_2 - m_1)g}{m_1 + m_2} = \frac{g}{5} \quad \text{and}$$

$$\frac{m_2 - m_1}{m_1 + m_2} = \frac{1}{5}$$

$$\frac{\frac{m_2}{m_1} - 1}{1 + \frac{m_2}{m_1}} = \frac{1}{5} \quad \text{let } \alpha = \frac{m_2}{m_1}$$

$$\frac{\alpha - 1}{\alpha + 1} = \frac{1}{5} \quad 5(\alpha - 1) = 5\alpha - 5 = \alpha + 1 \quad 4\alpha = 6$$

$$\alpha = \frac{3}{2}$$

